

## **A Novel Way to Prepare Soluble Graphene through Organic Functionalization on Graphene**

**Xiaoyan Zhang**<sup>1,2</sup>, Wesley R. Browne<sup>1</sup>, Bart J. van Wees<sup>2</sup> and Ben L. Feringa<sup>1</sup>

<sup>1</sup>Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

<sup>2</sup>Physics of Nanodevices, Zernike Institute for Advanced Materials, University of Groningen, The Netherlands

[xiaoyan.zhang@rug.nl](mailto:xiaoyan.zhang@rug.nl)

Soluble functionalized graphene was prepared through organic functionalization on graphene, which is confirmed by several techniques. And the obtained graphene is useful in several applications.

### **References**

- (1) X. Y. Zhang, A. C. Coleman, N. Katsonis, W. R. Browne, B. J. van Wees, B. L. Feringa, *Chem. Commun.*, **46** (2010), 7539.
- (2) X. Y. Zhang, L. L. Hou, A. Clossen, A. C. Coleman, O. Ivashenko, P. Rudolf, B. J. van Wees, W. R. Browne, B. L. Feringa, *Chemistry-A European Journal*, **17** (2011), 8957.
- (3) W. F. van Dorp, X. Y. Zhang, B. L. Feringa, J. B. Wagner, T. W. Hansen and J. Th. M. De Hosson, *Nanotechnology*, **22** (2011), 505303.
- (4) Y. Zhu, S. Murali, W. Cai, X. Li, J. Suk, J. Potts and R. Ruoff, *Adv. Mater.* **22** (2010), 3906.